

Latest Results on Orbitally Excited Bottom Mesons with CDF II Detector



Igor Gorelov

University of New Mexico

On behalf of

CDF Collaboration, F.N.A.L.



XXXIII International Conference on High Energy Physics

ICHEP 2006

July 26 -August 2, 2006, Moscow, Russia

1 – Outline

- Introduction.
- b -Physics at Tevatron with CDF II detector.
- CDF II Triggers and Datasets involved.
- Strange Orbital B_{sJ} Mesons with CDF II Detector.
 - ✓ Modes.
 - ✓ Predictions and Status.
 - ✓ Preparation of Candidates.
 - ✓ Selection of Candidates.
 - ✓ Results.
- Summary.

2 – Introduction

⇒ **H.Q.E.T. models:** $m_Q \rightarrow \infty, m_Q \gg \Lambda_{QCD}$

⇒ In HQET Bottom B -mesons $\bar{q}Q$ can be treated as a quark “hydrogen atom”:

- heavy bottom quark spin $s_Q = \frac{1}{2}^+$
decouples from light quark degrees of freedoms.
- the light anti-quark spin-parity $s_{\bar{q}} = \frac{1}{2}^-$ couples with its orbital momentum L .
- $\vec{j}_q = \vec{s}_{\bar{q}} + \vec{L}$ is a good quantum number.
- $\vec{J} = \vec{j}_q + \vec{s}_Q$ is a total momentum.


⇒ Consider L=1, B-mesons:

- $J = \frac{1}{2}^+ \oplus \frac{1}{2}^+$ or
- **B'_1 meson, $1^+ ({}^1P_1)$ and B'_0 meson, $0^+ ({}^1P_0)$.**
- $J = \frac{3}{2}^+ \oplus \frac{1}{2}^+$ or
- **B_1 meson, $1^+ ({}^1P_1)$ and B_2^* meson, $2^+ ({}^1P_2)$.**
- As m_Q is large, **but not infinite**, $\vec{j}_q - \vec{s}_Q$ interaction splits the mass states within these doublets.

3 – *b*- Physics at Tevatron@1.96 TeV

- Compare $\sigma(b\bar{b})$
 - $\Upsilon(4S) \approx 1 \text{ nb}$ (only B^0, B^+)
 - $Z^0 \approx 7 \text{ nb}$
 - $p\bar{p} \approx 20 \mu\text{b}$ ($|\eta| < 1.0$),
@1.96 TeV
- A reach variety of heavy quark and excited states are available at Tevatron
 - ... among them
 $B_s, B_c, B_J, \Lambda_b^0$ etc.
- enormous inelastic total cross-section of $\sim 60 \text{ mb}$
- *b*- Hadron Triggers required
 - based on leptons
 - based on displaced tracks
- high occupancy
- precise tracking with good resolution are required

4 – CDF II Detector at Tevatron





⇒ Critical for present results: COT (central tracker), SVX II (Si), μ -chambers

5 – *b*- Physics Triggers in CDF II


- ⇒ Trigger on Hadron Modes
- ⇒ of Heavy B-, D- Mesons, Baryons

- Level 1 eXtremely Fast Tracker (XFT)
Trigger: high p_T
- Level 2 Silicon Vertex Trigger (SVT):
large track d_0
- impact parameter resolution by SVX II
 $\sigma(d_0) = \sigma_{\text{beam}} \oplus \sigma_{\text{SVX}} = 47 \mu\text{m}$
- At a Level 3: a full event reconstruction.
- Improves Run I sensitivity by 4-5 orders of magnitude.
- **Present orbital *B*-mesons analysis**
use as a base mode:

$$B^+ \rightarrow \bar{D}^0 \pi^+, \bar{D}^0 \rightarrow K^+ \pi^-$$




⇒ J/ψ Trigger



- Reach of $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ mode
- Level 1 eXtremely Fast Tracker (XFT) Trigger: tracks are reconstructed in the COT and matched to hits in the Muon Chambers.
- Level 2: transparent
- Level 3: full reconstruction with cut $M(\mu^+ \mu^-) \in [2.7, 4.0] \text{ GeV}/c^2$

6 – Orbitally Excited B_{sJ} Mesons: Predictions

$\Rightarrow S-$ and $P-$ wave multiplets of strange B_s -mesons, similar to $B_{u,d}$



$\Rightarrow \dots$ and their possible decay modes:

- B_{sJ} states are isotop-singlets, $I=0$, **NO $B_s\pi$ modes allowed.**
- P -parity and $I-, S-$ are conserved.
- $\frac{1}{2}^+ \rightarrow \frac{1}{2}^- + 0^-, K$ in *S-wave*
- broad states
- $B'_{s1} \rightarrow B^+ K^-$, $1^+ \rightarrow 1^- + 0^-$
- $B_{s0}^* \rightarrow B^{*+} K^-$, $0^+ \rightarrow 0^- + 0^-$
- $\frac{3}{2}^+ \rightarrow \frac{1}{2}^- + 0^-, K$ in *D-wave*
- narrow states
- $B_{s1} \rightarrow B^{*+} K^-$, $1^+ \rightarrow 1^- + 0^-$
- $B_{s2}^* \rightarrow B^+ K^-$, $2^+ \rightarrow 0^- + 0^-$
- $B_{s2}^* \rightarrow B^{*+} K^-$, $2^+ \rightarrow 1^- + 0^-$ with $B^* \rightarrow B\gamma$, and γ is undetected.


⇒ Mass and Width predictions B_{sJ} states from several authors ...

State/ref.	Ebert <i>et al.</i>		Godfrey <i>et al.</i>		Eichten <i>et al.</i>		Falk <i>et al.</i>	
[MeV/ c^2]	M	Γ	M	Γ	M	Γ	M	Γ
B_{s0}^*	5841.0	-	5830.0	170	-	-	-	-
B'_{s1}	5859.0	-	5860.0	-	-	-	-	-
B_{s1}	5831.0	-	5860.0	-	5834.0	<1.0	5886.0	2.8 ± 1.2
B_{s2}^*	5844.0	-	5888.0	2.6	5846.0	1.0	5899.0	7 ± 3

- From HQET $\mathcal{B}(B_{s2}^* \rightarrow B^+ K^-)/\mathcal{B}(B_{s2}^* \rightarrow B^{*+} K^-) \sim 12.0$
- $B^{*+} \rightarrow B^+ \gamma$ with γ going undetected causes a shift of possible B_{s2}^*, B_{s1} peak positions by $\Delta M(B^{*+} - B^+) = 45.78 \text{ MeV}/c^2$ (see PDG)
- $B_{s2}^* \rightarrow B^+ K^-$ is expected to appear at $\mathbf{Q} \sim (70...130) \text{ MeV}/c^2$
- If B_{s2}^* is above $B^{*+} K^-$ threshold, a weaker signal by $45.78 \text{ MeV}/c^2$ less.
- $B_{s1} \rightarrow B^{*+} K^-$, a narrow peak at $\mathbf{Q} \sim (10...70) \text{ MeV}/c^2$, shifted due to γ .
- **Search for narrow signals in $\mathbf{Q} = M(B^+ K^-) - M(B^+) - M(K^-)$**

7 – Orbitally Excited B_{sJ} Mesons: Experimental Status

- OPAL (LEP): 1-st observation.
 149 ± 30 evts in $M(B^+ K^-)$.
 $M(B^+ K^-) = 5853 \pm 15 \text{ MeV}/c^2$.
 $B^+ K^-$ vs $B^{*+} K^-$, coarse resolution.
- DELPHI (LEP): a signal in
 $Q = M(B^+ K^-) - M(B^+) - M(K^-)$.
 $134 \pm 32(\text{stat}) \pm 12(\text{syst})$ evts
at $Q = 79.0 \pm 4.0(\text{stat}) \pm 2.0(\text{syst}) \text{ MeV}/c^2$.
Claimed $B_{s2}^* \rightarrow B^+ K^-$ observation.



DØ Collab. reported (Note 5027-Conf)

$135 \pm 31(\text{stat})$ evts
at $Q = 66.4 \pm 1.4(\text{stat}) \pm 2.0(\text{syst}) \text{ MeV}/c^2$,
 $\sigma(Q) = 4.7 \pm 1.2(\text{stat}) \text{ MeV}/c^2$
interpreted as a $B_{s2}^* \rightarrow B^+ K^-$
with $M = 5839.1 \pm 1.4(\text{stat}) \text{ MeV}$

8 – Orbitally Excited B_{sJ} Mesons: Candidate Preparation

⇒ Reconstruction of Candidates:

$$\begin{aligned} B_{sJ}^0 &\rightarrow B^+ K^-, \quad B^+ \rightarrow J/\psi K^+ \\ J/\psi &\rightarrow \mu^+ \mu^- + \text{chrg.conj.combs.} \end{aligned}$$

- $\mathcal{L} = 1066 \pm 64 \text{ pb}^{-1}$ with J/ψ trigger
- " J/ψ "($\mu^+ \mu^-$) $\in (2.9, 3.3) \text{ GeV}$
- 3-d Vertex Fit for " J/ψ "
- " J/ψ " $K^+ \in (4.6, 6.8) \text{ GeV}/c^2$
- 3-d Vertex Fit for " J/ψ " K^+
- " B^+ " K^- , B_{sJ}^0 candidate prepared.

⇒ Reconstruction of Candidates:

$$\begin{aligned} B_{sJ}^0 &\rightarrow B^+ K^-, \quad B^+ \rightarrow \bar{D}^0 \pi^+ \\ \bar{D}^0 &\rightarrow K^+ \pi^- + \text{chrg.conj.combs.} \end{aligned}$$

- $\mathcal{L} = 976 \pm 59 \text{ pb}^{-1}$ with 2-trk trigger
- " \bar{D}^0 "($K^+ \pi^-$) $\in (1.77, 1.97) \text{ GeV}$
- 3-d Vertex Fit for " \bar{D}^0 "
- " \bar{D}^0 " $\pi^+ \in (4.4, 6.6) \text{ GeV}/c^2$
- 3-d Vertex Fit for " B^+ "
- 4-track $B_{sJ}^0 \rightarrow "B^+" K^-$ candidate prepared.



9 – Orbitally Excited B_{sJ} Mesons: Candidate Selection

$$\Rightarrow B_{sJ}^0 \rightarrow B^+ K^-, B^+ \rightarrow J/\psi K^+$$

$$\Rightarrow B_{sJ}^0 \rightarrow B^+ K^-, B^+ \rightarrow \bar{D}^0 \pi^+$$

- For further selection of reconstructed candidates apply *Neural Network* software *NeuroBayes*[©], developed by Karlsruhe Univ. (Germany) team of CDF Collab.
 - Feed the topological quantities: L_{xy} , d_0 , $\text{Prob}(B^+)$ etc.
 - Feed the kinematic variables: p_T , $\cos(\theta^*)(K^-)$ etc.
 - Use **PID**(K^-) based on *ToF* or/and dE/dx
 - NN output from selected B^+ candidate for B_{sJ} selection.
- NN training:
 - **for background patterns:** exp. data from side bands of parent candidates
 - **for signal patterns:** MC Generator + CDF full simulation with **Q** shape generated as a background one.

⇒ The Neural Network produces the output distributed as:



⇒ The optimization of NN cut for every mode:


$$\text{F.o.M.} = N_{\text{cands}}^{\text{MC}}(NN) / \sqrt{N_{\text{cands}}^{\text{Data}}(NN)}$$

10 – B_{sJ} Mesons: Results


⇒ Make final cuts:

- the output of the \mathcal{NN}
 - $\mathcal{NN} > 0.5$ for $B^+ \rightarrow J/\psi K^+$
 - $\mathcal{NN} > 0.3$ for $B^+ \rightarrow \bar{D}^0 \pi^+$
- a number of candidates " B_{sJ} " → $B^+ K^-$ per event
 - $n_{cands}("B_{sJ}") < 4$
 - ... to further suppress a background.
- control background with wrong charge candidates " B_{sJ} " → $B^+ K^+$
- $Q = M("B^+ K^-") - M("B^+") - M(K^-)$

⇒ Experimental data: Q Spectra, two basic modes $B^+ \rightarrow J/\psi K^+$, $\bar{D}^0 \pi^+$



⇒ Both modes are added.



⇒ Two narrow peaks at


⇒ $Q \sim 10, \sim 70 \text{ MeV}/c^2$ are visible.

⇒ Wrong charge $B^+ K^+$ are shown.

$\Rightarrow B_{sJ} \rightarrow B^+ K^-$, ($B^+ \rightarrow J/\psi K^+$)

\Rightarrow Fit Results.

State	$B_{s1} \rightarrow B^{*+} K^-$
B^+ Mode	$B^+ \rightarrow J/\psi K^+$
Q [MeV]	10.87 ± 0.19
σ [MeV]	0.64 ± 0.25
N_{evts}	16.98 ± 5.14
State	$B_{s2}^* \rightarrow B^+ K^-$
B^+ Mode	$B^+ \rightarrow J/\psi K^+$
Q [MeV]	67.03 ± 0.44
σ [MeV]	1.79 ± 0.42
N_{evts}	44.15 ± 13.36



\Rightarrow unbinned N.L.L. fit

\Rightarrow Assign narrow peaks


\Rightarrow @ $Q \sim 10.9$ MeV $\mapsto B_{s1}$

\Rightarrow @ $Q \sim 67.0$ MeV $\mapsto B_{s2}^*$

$\Rightarrow B_{sJ} \rightarrow B^+ K^-$, ($B^+ \rightarrow \bar{D}^0 \pi^+$)

\Rightarrow Fit Results.

State	$B_{s1} \rightarrow B^{*+} K^-$
B^+ Mode	$B^+ \rightarrow \bar{D}^0 \pi^+$
Q [MeV]	10.68 ± 0.46
σ [MeV]	1.18 ± 0.56
N_{evts}	20.66 ± 7.12
State	$B_{s2}^* \rightarrow B^+ K^-$
B^+ Mode	$B^+ \rightarrow \bar{D}^0 \pi^+$
Q [MeV]	66.85 ± 0.76
σ [MeV]	2.88 ± 0.75
N_{evts}	55.74 ± 19.20



\Rightarrow unbinned N.L.L. fit

\Rightarrow Assign narrow peaks

\Rightarrow @ $Q \sim 10.9$ MeV $\mapsto B_{s1}$


\Rightarrow @ $Q \sim 67.0$ MeV $\mapsto B_{s2}^*$

⇒ Fit Results for combined statistics:

- $Q(B_{s1}) = 10.73 \pm 0.21(\text{stat}) \text{ MeV}$
 $N_{\text{evts}} = 36.4 \pm 9.0(\text{stat})$
- $Q(B_{s2}^*) = 66.96 \pm 0.39(\text{stat}) \text{ MeV}$
 $N_{\text{evts}} = 94.8 \pm 23.4(\text{stat})$
- From unbinned N.L.L. fits,
 assuming
 $-2 \ln \mathcal{L}/\mathcal{L}_0 \sim \chi^2$
 - B_{s1} significance is 6.3σ
 - B_{s2}^* significance is 7.7σ

$$\Rightarrow B_{sJ} \rightarrow B^+ K^-$$

$$\Rightarrow (B^+ \rightarrow J/\psi K^+, \bar{D}^0 \pi^+)$$




⇒ unbinned N.L.L. fit

⇒ Significance of B_{s1} with toy MC

- B_{s2}^{*} has already been seen
- B_{s1} is a **first observation**
- Search window
- $\mathbf{Q} \in (0.0, 50) \text{ MeV}$
- Generate only background and B_{s2}^{*} and fit it
- Figure of Merit: $-2 \ln \mathcal{L}/\mathcal{L}_0$
- p-value for B_{s1} is $7.32 \cdot 10^{-6}$

⇒ p-value



⇒ $-2 \ln \mathcal{L}/\mathcal{L}_0$ from unbinned N.L.L. fit


11 – Summary

- $Q(B_{s1}) = 10.73 \pm 0.21(\text{stat}) \pm 0.14(\text{syst}) \text{ MeV}$
 $N_{evts} = 36.4 \pm 9.0(\text{stat}), 6.3\sigma$
p-value for B_{s1} is $7.32 \cdot 10^{-6}$, $Q \in (0, 50) \text{ MeV}$
The presented results are a first observation of a narrow B_{s1} state.
- $Q(B_{s2}^*) = 66.96 \pm 0.39(\text{stat}) \pm 0.14(\text{syst}) \text{ MeV}$
 $N_{evts} = 94.8 \pm 23.4(\text{stat}), 7.7\sigma$
- $M(B_{s1}) = 5829.41 \pm 0.21(\text{stat}) \pm 0.14(\text{syst}) \pm 0.6(\text{PDG}) \text{ MeV}$
- $M(B_{s2}^*) = 5839.64 \pm 0.39(\text{stat}) \pm 0.14(\text{syst}) \pm 0.5(\text{PDG}) \text{ MeV}$
- $\Delta M = 10.20 \pm 0.44(\text{stat}) \pm 0.35(\text{PDG}) \text{ MeV}$
- **The widths of measured B_{sJ} states are found to be at the level of a detector resolution.**
- **CDF II measurements on B_{s2}^* are in agreement with other experiments.**

THE END OF THE TALK.

12 – Backup Slides


⇒ ***S*– and *P*–wave multiplets of non-strange $B_{u,d}$ -mesons:**



⇒ ... and their possible decay modes:

- **strong decays** with emission of $\pi, 0^-$
 - P –parity and I – are conserved.
- $\frac{1}{2}^+ \rightarrow \frac{1}{2}^- + 0^-$,
 π emitted in ***S*–wave, $L = 0$** .
 - $\Gamma_{theor.} \sim 100 \text{ MeV}/c^2$, broad.
 - $B'_1 \rightarrow B^* \pi$, $1^+ \rightarrow 1^- + 0^-$
 - $B_0^* \rightarrow B \pi$, $0^+ \rightarrow 0^- + 0^-$
- $\frac{3}{2}^+ \rightarrow \frac{1}{2}^- + 0^-$,
 π emitted in ***D*–wave, $L = 2$** .
 - $\Gamma_{theor.} \sim 10 - 20 \text{ MeV}/c^2$, narrow.
 - $B_1 \rightarrow B^* \pi$, $1^+ \rightarrow 1^- + 0^-$
 - $B_2^* \rightarrow B \pi$, $2^+ \rightarrow 0^- + 0^-$
 - $B_2^* \rightarrow B^* \pi$, $2^+ \rightarrow 1^- + 0^-$
 with $B^* \rightarrow B\gamma$, and γ is undetected.

⇒ Earlier (Fall 2005) CDF presented results on orbitally excited B_1 and B_2^* mesons.



⇒ using two main CDF B-Physics Trigger streams.

- $B^+ \rightarrow J/\psi K^+$, J/ψ Trigger.
- $B^+ \rightarrow \bar{D}^0 \pi^+$, “2-trk Trigger”.
- $Q = M(B^+ \pi^-) - M(B^+) - M(\pi^-)$
- $M(B_1) = 5734 \pm 3(\text{stat}) \pm 2(\text{syst}) \text{ MeV}/c^2$
- $M(B_2^*) = 5738 \pm 5(\text{stat}) \pm 1(\text{syst}) \text{ MeV}/c^2$
- DØ has presented recently (DØ Note 5026-CONF) the measurements using a total available statistics of 1 fb^{-1}

⇒ Quantities used to feed NN

$B_{sJ}^0 \rightarrow B^+ K^-$, $B^+ \rightarrow J/\psi K^+$

- $B^+ \rightarrow J/\psi K^+$
 - $L_{xy}(J/\psi)$ w.r.t. PVX
 - $p_T(K^+)$
 - $d_0(K^+)$ and $d_0(K^+)/\sigma$
 - $d_0(B^+)$
 - Prob(B^+)
- $B_{sJ} \rightarrow B^+ K^-$
 - $d_0(B_{sJ})/\sigma$
 - \mathcal{NN} output from B^+ candidate
 - **PID**(K^-) based on *ToF*
 - $\cos(\theta^*)(K^-)$ in B_{sJ} rest frame
 - $\eta(K^-)$, pseudorapidity
 - **NN training:** bgr., exp. data (s.b.)...

$B_{sJ}^0 \rightarrow B^+ K^-$, $B^+ \rightarrow \bar{D}^0 \pi^+$

- $B^+ \rightarrow \bar{D}^0 \pi^+$
 - $L_{xy}(B^+)/\sigma$ w.r.t. PVX
 - $p_T(\pi^+)$
 - $d_0(B^+)$
 - Prob(B^+)
- $B_{sJ} \rightarrow B^+ K^-$
 - $d_0(B_{sJ})$
 - $p_T(B_{sJ})$
 - \mathcal{NN} output from B^+ candidate
 - **PID**(K^-) based on *ToF* or dE/dx
 - **M**(B^+), inv. mass
 - ... and signal, **BGEN + CDF sim.**

$\Rightarrow B^+ \rightarrow J/\psi K^+, \bar{D}^0 \pi^+$ after optimized NN Selection...

